



Division: Pharmacy Policy	Subject: State of Florida's Agency for Health Care Administration's Prior Authorization Criteria
Original Development Date: Original Effective Date: Revision Date:	April 20, 2009 April 24, 2012; September 28, 2012; April 29, 2013, February 20, 2018, July 15, 2020, September 22, 2020, March 11, 2022

## **H.P. Acthar Gel (repository corticotropin injection) and Purified Cortrophin Gel (corticotropin injection)**

**LENGTH OF AUTHORIZATION:** Per Titration Schedule (length of approval shall not exceed the quantity (mL) required to properly treat and taper)

### **REVIEW CRITERIA:**

#### **Collagen disease (Approve for 3 months):**

- Patient must be  $\geq 2$  years old.
- Treatment of acute exacerbations or maintenance therapy.
- Must have a diagnosis of a collagen disorder (e.g. systemic lupus erythematosus and systemic dermatomyositis (polymyositis).
- If the diagnosis is systemic lupus erythematosus, patient has had a trial of 1 agent from each of the following, and experienced inadequate response or intolerance:
  - a) Corticosteroid: prednisone, intravenous dexamethasone
  - b) Hydroxychloroquine, azathioprine, cyclosporine, cyclophosphamide, or mycophenolate
  - c) Intravenous methylprednisolone
- For other diagnoses of collagen disorder, patient must have failed corticosteroid therapy (e.g. dexamethasone, hydrocortisone, methylprednisolone, prednisone...).
- Medication must be prescribed by rheumatologist or a specialist in this field of study.

#### **Disorder of eye (Approve for 3 months):**

- Patient must be  $\geq 2$  years old.
- Use for severe acute and chronic allergic and inflammatory processes involving the eye and its adnexa such as: keratitis, iritis, iridocyclitis, diffuse posterior uveitis and choroiditis, optic neuritis, chorioretinitis, and anterior segment inflammation.
- Patient must have failed corticosteroid therapy (e.g. dexamethasone, hydrocortisone, methylprednisolone, prednisone . . .).
- Medication must be prescribed by ophthalmologist or a specialist in this field of study.

#### **Disorder of skin (Approve for 3 months):**

- Patient must be  $\geq 2$  years old.
- For the treatment of dermatologic diseases such as erythema multiforme and Stevens-Johnson syndrome.

#### **Infantile Spasms (Approve for 3 months):**

- Patient must be less than 2 years old.
- Must have a diagnosis of **West Syndrome (infantile spasms)** verified by progress notes, discharge notes, or health conditions.
- Medication must be prescribed by neurologist or a specialist in this field of study.

**Acute Exacerbations in Adults with Multiple Sclerosis (MS) (Approved for one month):**

- Patient must be  $\geq 18$  years old.
- Must have a diagnosis of acute exacerbation of MS verified by progress notes, discharge notes, or health conditions.
- Failure of a recent (within the last 30 days) trial of at least 3 day course of corticosteroid therapy (e.g. dexamethasone, hydrocortisone, methylprednisolone, prednisone . . .). for acute exacerbations of MS, unless contraindicated or clinically significant adverse effects are experienced.
- Patient is adherent to disease modifying treatment (e.g. Aubagio, Avonex Betaseron, Copaxone, Gilenya, Rebif, Tecfidera.).
- Medication must be prescribed by neurologist or a specialist in this field of study.

**Acute Exacerbations of Inflammatory disorder of musculoskeletal system (Approve for 3 months):**

- Patient must be  $\geq 2$  years old.
- Must be diagnosed with a musculoskeletal disease (e.g. psoriatic arthritis, rheumatoid arthritis, juvenile rheumatoid arthritis, or ankylosing spondylitis) verified by progress notes, discharge notes, or health conditions.
- Treatment is for acute exacerbations, short term adjunct therapy.
- Medication must be prescribed by rheumatologist or a specialist in this field of study.
- Patient must have failed corticosteroid therapy (e.g. dexamethasone, hydrocortisone, methylprednisolone, prednisone....).

**Nephrotic syndrome (Approve for 3 months):**

- Patient must be  $\geq 2$  years old.
- Must have a diagnosis of nephrotic syndrome, idiopathic, without uremia or due to lupus erythematosus verified by progress notes, discharge notes, or health conditions.
- Medication must be prescribed by nephrologist or a specialist in this field of study.
- Patient must have failed corticosteroid therapy (e.g. dexamethasone, hydrocortisone, methylprednisolone, prednisone....).

**Sarcoidosis (Approve for 3 months):**

- Patient must be  $\geq 2$  years old.
- Must have the diagnosis of sarcoidosis verified by progress notes, discharge notes, or health conditions.
- Medication must be prescribed by rheumatologist or a specialist in this field of study.
- Patient must have failed corticosteroid therapy (e.g. dexamethasone, hydrocortisone, methylprednisolone, prednisone....).

**Transfusion reaction due to serum protein reaction (Approve for 3 months):**

- Patient must be  $\geq 2$  years old.
- Must have a diagnosis of serum sickness.

**CONTINUATION OF THERAPY (UP TO 3 MONTHS):**

Non-acute indications:

- Patient met initial review requirements
- Positive response to therapy submitted (supporting documentation required)
- Dosage and administration does not exceed FDA approved maximum for the patient's indication.
- Supporting documentation if dose requested exceeds FDA approved maximum.

**DOSING AND ADMINISTRATION:**

- Refer to product labeling at <https://www.accessdata.fda.gov/scripts/cder/daf/>