

Division: Pharmacy Policy	Subject: State of Florida's Agency for Health Care Administration's
	Prior Authorization Criteria
Original Development Date: Original Effective Date:	May 6, 2022
Revision Date:	

# **Intravenous and Injectable Iron Agents**

Preferred: Ferrlecit®, Sodium Ferric Gluconate Complex

Non-Preferred: Feraheme®, Ferumoxytol, Infed®, Injectafer®, Monoferric®, Triferic®, Venofer® (refer to specific

criteria)

**LENGTH OF AUTHORIZATION**: Initial Therapy - 3 months

Continuation of therapy - 6 months

## **REVIEW CRITERIA**:

- Medication requested must have a documented FDA approved indication and the patient must be within the FDA approved age limits.
- Patient is intolerant or had an unsatisfactory response to oral iron.
- Patient must have serum ferritin  $\leq$  300 ng/ml and transferrin saturation  $\leq$  20% (baseline lab data drawn within 30 days of PA submission must be provided).
- Patients with Chronic Kidney Disease (CKD) on erythropoiesis stimulating therapy must have serum ferritin ≤ 500 ng/ml and transferrin saturation ≤ 30% (baseline lab data drawn within 30 days of PA submission must be provided.
- The patient must have documented trial and failure on medications on the Preferred Drug List (PDL) or there is a reason (allergy, contraindication) that preferred drugs cannot be used.
- The requested medication's corresponding generic (if available) has been attempted and failed.

## **CONTINUATION OF THERAPY:**

- Patient has met initial review criteria; AND
- Documentation of improved clinical response compared to baseline labs (supporting documentation and official lab results required); **AND**
- Patient has not experienced any treatment-restricting adverse effects; AND
- Dosing is appropriate as per labeling or is supported by compendia.

### Venofer (iron sucrose):

#### **Adult Patients**

## **Initial Review Therapy**

- Patient must be  $\geq 18$  years of age.
- Patient must have a documented diagnosis of iron deficiency anemia AND one of the following:
  - Hemodialysis Dependent CKD
  - Non-Dialysis Dependent CKD
  - o Peritoneal Dialysis Dependent CKD



Division: Pharmacy Policy	Subject: State of Florida's Agency for Health Care Administration's
	Prior Authorization Criteria
Original Development Date: Original Effective Date:	May 6, 2022
Revision Date:	

- The patient must have documented trial and failure on medications on the PDL or there is a reason (allergy, contraindication) that preferred drugs cannot be used.
- Patient must have serum ferritin ≤ 300 ng/ml and transferrin saturation ≤ 20% (lab data drawn within 30 days of PA submission must be provided).
- Patients with CKD on erythropoiesis stimulating therapy must have serum ferritin  $\leq$  500 ng/ml and transferrin saturation  $\leq$  30% (lab data drawn within 30 days of PA submission must be provided)

#### **Pediatric Patients**

#### **Initial Review Criteria**

- Patient must be  $\geq 2$  years of age.
- Patient must have a documented diagnosis of iron deficiency anemia AND one of the following:
  - o Hemodialysis Dependent CKD
  - o Non-Dialysis Dependent CKD
  - Peritoneal Dialysis Dependent CKD receiving concurrent erythropoiesis stimulating therapy
- The patient must have documented trial and failure on medications on the PDL or there is a reason (allergy, contraindication) that preferred drugs cannot be used.
- Patient must have serum ferritin ≤ 300 ng/ml and transferrin saturation ≤ 20% for Hemodialysis Dependent CKD and Non-Dialysis Dependent CKD (baseline lab data drawn within 30 days of PA submission must be provided).
- Patient must have serum ferritin  $\leq 500$  ng/ml and transferrin saturation  $\leq 30\%$  for Peritoneal Dependent CKD (baseline lab data drawn within 30 days of PA submission must be provided).

#### **CONTINUATION OF THERAPY:**

- Patient meets initial review criteria; AND
- Documentation of improved clinical response compared to baseline labs (supporting documentation and official lab results required); **AND**
- Patient has not experienced any treatment-restricting adverse effects; AND
- Dosing is appropriate as per labeling or is supported by compendia.

## DOSING AND ADMINISTRATION:

Refer to product labeling at <a href="https://www.accessdata.fda.gov/scripts/cder/daf/">https://www.accessdata.fda.gov/scripts/cder/daf/</a>