



Division: Pharmacy Policy	Subject: State of Florida's Agency for Health Care Administration's Prior Authorization Criteria
Original Development Date: Original Effective Date: Revision Date:	April 30, 2021

VERQUVO™ (vericiguat)

LENGTH OF AUTHORIZATION: Up to one year

REVIEW CRITERIA:

- Patient must be ≥ 18 years of age.
- Patient must have a diagnosis of symptomatic chronic heart failure (New York Heart Association [NYHA] class II-IV).
- Patient must have a left ventricular ejection fraction (LVEF) less than 45%.
- Documentation of recent (within 6 months) hospitalization due to CHF or a demonstrated need for outpatient IV diuretics (within 3 months).
- Documentation of prior or current therapy with an ACEI/ARB and/or beta-blocker and/or mineralocorticoid receptor antagonist (MRA).
- Women of childbearing potential should have a negative pregnancy test collected prior to therapy initiation.
- Patient may not use with another soluble guanylate cyclase (sGC) stimulator or a phosphodiesterase-5 (PDE-5) inhibitor.
- Must be prescribed by or in consultation with a cardiologist.

CONTINUATION OF THERAPY:

- Patient met initial review criteria.
- Documentation of therapy compliance.
- Patient has experienced a positive clinical response to Verquvo (e.g. decreased hospitalizations).

DOSING AND ADMINISTRATION:

- Dosage form: 2.5 mg, 5 mg and 10 mg tablets
- Recommended dose: 2.5 mg by mouth once daily titrated to maintenance dose of 10 mg daily