

| Division: Pharmacy Policy | Subject: State of Florida's Agency for Health Care Administration's |
|----------------------------|---|
| | Prior Authorization Criteria |
| Original Development Date: | January 2, 2020 |
| Original Effective Date: | |
| Revision Date: | March 9, 2021, October 4, 2021, March 31, 2022 |
| | |

VYONDYS 53 (golodirsen)

LENGTH OF AUTHORIZATION: SIX MONTHS

REVIEW CRITERIA:

- Patient must have the diagnosis of Duchenne Muscular Dystrophy (DMD).
- Submission of medical records including the following:
 - Genetic testing confirming the patient has a mutation of the DMD gene that is amenable to exon 53 skipping.
 - o Baseline renal function tests (i.e., glomerular filtration rate GFR).
- Medication is prescribed by or in consultation with a neurologist or a physician who specializes in treatment of DMD (i.e., pediatric neurologist, cardiologist or pulmonary specialist).
- Patient has been on stable dose of oral corticosteroids for at least 24 weeks prior to starting therapy unless contraindicated or intolerant.
- Patient is not concurrently treated with other DMD antisense oligonucleotides (e.g., Casimersen, Viltolarsen, or Eteplirsen).
- If the patient is ambulatory, functional level determination of baseline assessment of ambulatory function (six-minute walk test (6MWT), time to run/walk 10-meter test (TTRW), time to climb 4-stair test (TTCLIMB), time to stand (TTSTAND) or North Star Ambulatory Assessment (NSAA)) is required.
- If not ambulatory, patient must have a Brooke Upper Extremity Function Scale of five or less documented OR a Forced Vital Capacity of 30% or more.

CONTINUATION OF THERAPY:

- Patient met initial review criteria.
- Documentation of improvement, maintenance, or slowing of disease progression:
 - o For ambulatory patients- Submission of 6MWT, TTRW, TTCLIMB, TTSTAND or NSAA.
 - o For non-ambulatory patients- Submission of Brooke Upper Extremity Function Scale (five or less) documented OR a Forced Vital Capacity documented (30% or more).
- Submission of renal function tests (i.e., GFR).
- Patient is not concurrently treated with other DMD antisense oligonucleotides (e.g., Casimersen, Viltolarsen, or Eteplirsen).

DOSING AND ADMINISTRATION:

- Refer to product labeling at https://www.accessdata.fda.gov/scripts/cder/daf/
- Available as 100 mg/2 mL single-dose vial



| Division: Pharmacy Policy | Subject: State of Florida's Agency for Health Care Administration's |
|----------------------------|---|
| | Prior Authorization Criteria |
| Original Development Date: | January 2, 2020 |
| Original Effective Date: | |
| Revision Date: | March 9, 2021, October 4, 2021, March 31, 2022 |
| | |